



LITERATURA

1. Vyzula R a kol. Modrá kniha České onkologické společnosti. Brno: Masarykův onkologický ústav, 2018: 260 s.
2. Gurney H. How to calculate the dose of chemotherapy. British Journal of Cancer. 2002; 86(8): 1297–1302.
3. Sacco JJ, et al. The average body surface area of adult cancer patients in the UK: a multicentre retrospective study. PloS One. 2010; 5(1): e8933.
4. Mosteller RD. Simplified Calculation of Body-Surface Area. The New England Journal of Medicine. 1987; 317(17): 1098–1098.
5. Haycock GB, et al. Geometric method for measuring body surface area: A height-weight formula validated in infants, children, and adults. The Journal of Pediatrics. 1978; 93(1): 62–66.
6. Výpočet hmotností segmentů lidského těla. BIOMECH.FTVS.CUNI.CZ [online]. [cit. 2018-09-10] Dostupné z: http://biomech.ftvs.cuni.cz/pbpbk/kompendium/biomechanika/geometrie_hmotnost_vypocet.php.
7. Owen D, et al. Optimal chemotherapy dosing in a bilateral lower extremities amputee with metastatic pancreatic adenocarcinoma. Gastrointestinal Cancer Research: GCR. 2013; 6(3): 93.
8. Hutson PR. Using body surface area adjustments in amputees. American Journal of Hospital Pharmacy. 1985; 42(5): 1048.
9. Wallace AB. The exposure treatment of burns. The Lancet. 1951; 257(6653): 501–504.
10. Smith JJ, et al. A comparison of serial halving and the rule of nines as a pre-hospital assessment tool in burns. British Journal of Plastic Surgery. 2005; 58(7): 957–967.
11. Colangelo PM, et al. Two methods for estimating body surface area in adult amputees. American Journal of Hospital Pharmacy. 1984; 41(12): 2650.
12. O'Marcaigh, AS, et al. Variation in the practice of dose reduction of chemotherapeutic agents after weight loss or amputation. Journal of Pediatric hematology/oncology. 1995; 17(2): 172–175.
13. O'Marcaigh AS, et al. Dose reduction of chemotherapeutic agents after weight loss. American Journal of Clinical Oncology. 1997; 20(2): 193–195.