



Závěr

Pečlivá kontrola arteriální hypertenze je přínosná v primární i sekundární prevenci CMP. Správně vedenou prevencí aterosklerotických vaskulárních

onemocnění můžeme přispět ke snížení počtu příhod, zlepšení kvality a prodloužení života našich pacientů.

LITERATURA

1. Donkor ES. Stroke in the 21st Century: A Snapshot of the Burden, Epidemiology, and Quality of Life. *Stroke Res Treat.* 2018; 2018: 3238165. doi: 10.1155/2018/3238165.
2. Kotseva K, De Backer G, De Bacquer D. Lifestyle and impact on cardiovascular risk factor control in coronary patients across 27 countries: Results from the European Society of Cardiology ESC-EORP EUROASPIRE V registry. *European Journal of Preventive Cardiology* 2019; 26(8). doi: <https://doi.org/10.1177/2047487318825350>.
3. Lüscher TF. Epidemiology of cardiovascular disease: the new ESC Atlas and beyond. *European Heart Journal* 2018; 39(7): 489–492. doi: <https://doi.org/10.1093/eurheartj/ehy070>
4. Samsa GP, Bian J, Lipscomb J, Matchar DB. Epidemiology of recurrent cerebral infarction: a medicare claims-based comparison of first and recurrent strokes on 2-year survival and cost. *Stroke* 1999; 30: 338–349.
5. Feng W, Hendry HM, Adams RJ. Risk of recurrent stroke, myocardial infarction, or death in hospitalized stroke patients. *Neurology* 2010; 74(7). doi: <https://doi.org/10.1212/WNL.0b013e3181c7776>.
6. Zanchetti A, Liu L, Mancia G, et al. Blood pressure and low-density lipoprotein-cholesterol lowering for prevention of strokes and cognitive decline: a review of available trial evidence. *J Hypertens* 2014; 32: 1741–1750.
7. Cífková R, Wohlfahrt P, Krajčoviechová A, et al. Blood pressure control and risk profile in poststroke survivors: a comparison with the general population. *J Hypertens.* 2015; 33(10): 2107–2114. doi: 10.1097/HJH.0000000000000660. PMID: 26200884.
8. Hong SK. Blood Pressure Management for Stroke Prevention and in Acute Stroke. *J Stroke* 2017; 19(2): 152–165. doi: 10.5853/jos.2017.00164.
9. Kohok DD, Sico JJ, Bae F, et al. Post-stroke hypertension control and receipt of health care services among veterans. *JCH* 2018; 20(2): 382–387. doi: <https://doi.org/10.1111/jch.13194>
10. Ahmed N, Audebert H, Turc G, et al. Consensus statements and recommendations from the ESO-Karolinska Stroke Update Conference, Stockholm 11–13 November 2018. *European Stroke Journal* 2019; 4(4): 307–317. doi: <https://doi.org/10.1177/2396987319863606>.
11. Williams B, Mancia G, Spiering W, et al. 2018 ESC/ESH Guidelines for the management of arterial hypertension: The Task Force for the management of arterial hypertension of the European Society of Cardiology (ESC) and the European Society of Hypertension (ESH). *European Heart Journal* 2018; 39(33): 3021–3104. doi: <https://doi.org/10.1093/eurheartj/ehy339>.
12. Williamson JD, Supiano MA, Applegate WB, et al. SPRINT Research Group. Intensive vs standard blood pressure control and cardiovascular disease outcomes in adults aged ≥ 75 years: a randomized clinical trial. *JAMA* 2016; 315: 2673–2682.
13. Ettehad D, Emdin CA, Kiran A, et al. Blood pressure lowering for prevention of cardiovascular disease and death: a systematic review and meta-analysis. *Lancet* 2016; 387: 957–967.
14. Emdin CA, Rahimi K, Neal B, et al. Blood pressure lowering in type 2 diabetes: a systematic review and meta-analysis. *JAMA* 2015; 313: 603–615.
15. PROGRESS Collaborative Group Randomised trial of a perindopril-based blood-pressure-lowering regimen among 6,105 individuals with previous stroke or transient ischaemic attack. *Lancet.* 2001; 358(9287): 1033–1041.